



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

dehisce, and spores germinate in the closed sporangia. As a result, dwarf male prothallia are produced, antheridia sometimes appearing at the three- or four-celled stage. Free spores under the same conditions never produce such prothallia, not having developed sexual organs at the conclusion of the experiment. The dwarf males do not burst through the wall of the sporangium, and ultimately die. It was found also that both free spores and those inclosed in sporangia germinate in darkness. The suggested explanation of the formation of the dwarf male prothallia under the conditions described has no foundation in experimental work. It is to be hoped that the day of imaginary "explanations" is about over.—J. M. C.

**Periodicity of algae.**—BROWN<sup>29</sup> has studied the appearance and disappearance of algae in selected ponds and streams in the vicinity of Bloomington, Indiana. In this region, an alga growing under "steady normal conditions" remains in a healthy vegetative state throughout the year. A sudden change in external conditions checks this vegetative growth, and induces a resting stage or sexual reproduction. In reference to specific plants, *Spirogyra nitida* is the most abundant of the Conjugatae in the region studied, and *S. varians* is the most widely distributed alga, conjugating at all seasons of the year when exposed to hard conditions (as the drying-up of a pond); *Chaetophora* thrives best in slightly stagnant water at a temperature between 5° and 25° C.; *Draparnaldia* finds its most congenial conditions in flowing surface water between 1° and 15° C.—J. M. C.

**Cytology of Synchytrium.**—GRIGGS<sup>30</sup> has continued an investigation on *S. decipiens* begun by F. L. STEVENS, to whom he is indebted for material. There are 500 to 800 free nuclei in the cyst when cell walls begin to appear; but most of the study was upon cysts with 100 to 300 free nuclei. While no centrosomes were found in the metaphase and anaphase, in the telophase there are large asters with centrosomes at the center, whose origin has not yet been determined. As the nuclear vacuole forms about the chromosomes, the coarse rays of the aster bend about it and become transformed into the thick nuclear membrane characteristic of the genus. It is hoped that a further study will throw some light upon systematic relations.—CHARLES J. CHAMBERLAIN.

**Sporangia of Lycopodiaceae.**—As a result of her study of the sporangium-bearing organs of the Lycopodiaceae, Miss SYKES<sup>31</sup> has arranged the different species of *Lycopodium* in a continuous series based on the shape and structure of the sporophyll, the position of the sporangium, and the position of the line of dehiscence. The evidence adduced seems scarcely sufficient to warrant the con-

---

<sup>29</sup> BROWN, HARRY B., Algal periodicity in certain ponds and streams. Bull. Torr Bot. Club 35: 223-248. 1908.

<sup>30</sup> GRIGGS, ROBT. F., On the cytology of Synchytrium. Ohio Naturalist 8: 277-286. pl. 20. 1908.

<sup>31</sup> SYKES, M. G., Notes on the morphology of the sporangium-bearing organs of the Lycopodiaceae. New Phytol. 7: 41-60. pls. 2, 3. 1908.